Company Name	
Crop Name	Bajara
Variety	
Types Of Variety	
(Hybrid/Composite/Synthetic/Improved/Single Cross/Double	
Cross)	

S No	Characteristics	States	Note	Hybrid/Research Variety	Female	Male
1	2	3	4	5	6	7
1	Plant : Anthocyanin colorationof first leaf sheath	Absent	1			
		Present	9			
	Plant :Growth habit	Erect	1			
2		Intermediate	5			
		Spreading	7			
	Time of spike emergence (50% plants with at least one spike emerged fully)	Very early (<43 days)	1			
		Early (43-46 days)	3			
3		Medium (47-50)	5			
		Late (51-54 days)	7			
		Very late (>54 days)	9			
4	Land, Chardle and and	Absent	1			
4	Leaf: Sheath pubescence	Present	9			
	Leaf: Sheath length (cm)	Short (<11)	3			
5		Medium (11-15)	5			
		Long (>15)	7			
	+	Very short (<41)	1			
		Short (41-50)	3			
6	Leaf: Blade length (cm)	Medium (51-60)	5			
		Long (61-70)	7			
		Very long (>70)	9			
	-	Narrow (<3)	3			
7	Leaf: Blade width (at widest point) (cm)	Medium (3-4)	5			
,	Lear : Blade width (at widest point) (em)	Broad (>4)	7			
		Absent	1			
8	Spike:Stigma Pigmentatio	Present	9			
		Yellow	3			
9	Spike: Anther colour	Brown	5			
,		Purple	7			
		Absent	1			
10	Plant: Node pubescence		9			
		Present	3			
1.1	Dlant, Number of nodes	Low (<11)				
11	Plant: Number of nodes	Medium (11-15)	5 7			
		High (>15)				
	Plant: Node pigmentation	Whitish	1			
10		Green	2			
12		Brown	3			
		Red	4			
		Purple	5			
	Plant: Internode pigmentation (between 3 rd & 4 th node from top)	Whitish	1			
1.0		Green	2			
13		Brown	4			
		Red	6			
		Purple	7			
	Spike exsertion	Incomplete	1			
14		Partial	3			
		Complete	5			
	Spike: Length (cm)	Very small (< 11)	1			
		Small (11-20)	3			
15		Medium (21-30)	5			
		Long (31-40)	7			
		Very long (>40)	9			
1.0		Absent	1			
16	Spike : Anthocyanin pigmentation of glume	Present	9			
1-	0.11 D.1.1	Absent	1			
17	Spike: Bristle	Present	9			
					ı	

S No	Characteristics	States	Note	Hybrid/Research Variety	Female	Male
1	2	3	4	5	6	7
18	Spike: Bristle colour	Yellow	1			
		Green	2			
		Brown	3			
		Purple	5			
	Spike: Bristle Appearance	Non prominent (Bristle length	3			
19		< 2mm from the ear head)	J			
1)		Prominent (Bristle length >	5			
		2mm from the ear head)	3			
	Spike: Girth at maximum point (excluding bristles) (cm)	Thin (<1.6)	3			
20		Medium (1.6-3.0)	5			
		Thick (>3.0)	7			
	Plant: Number of productive tillers	Monoculm (<2)	1			
21		Low (2-3 tillers)	3			
21		Medium (4-6 tillers)	5			
		High (>6 tillers)	7			
		Very short (<101)	1			
		Short (101-150)	3			
22	Plant: Height (including spike) (cm)	Medium (151-200)	5			
		Tall (201-250)	7			
		Very tall (>250)	9			
	Spike shape:	Cylindrical	1			
		Conical	2			
		Spindle	3			
		Candle	4			
23		Lanceolate	5			
		Dumb-bell	6			
		Club	7			
		Oblanceolate	8			
		Globose	9			
	Spike: Tip sterility	Absent	1			
24		Present	9			
25	Spike: Density	Loose	3			
		Semi-compact	5			
		Compact	7			
	Seed: Colour	Whitish	1			
		Cream	2			
		Yellow	3			
26		Grey	4			
		Deep grey	5			
		Grey brown	6			
		Yellow brown	7			
		Obovate	3		<u> </u>	
27	Seed: Shape	Elliptical	5			
		Hexagonal	7			
		Globular	9			
		Very low (<5)	1			
28	Seed: Weight of 1000 grains (g)	Small (5.0-7.5)	3			
		Medium (7.6-10.0)	5			
		Bold (10.1-12.5)	7			
			9			
		Very bold (>12.5)	9		1	