| Company Name | | | | | | | | | |
|--------------|---|------------------------|------|----------------------------|--------|------|--|--|--|
| Crop Name | | Groundnut | | | | | | | |
| Variety | | | | | | | | | |
| Types C | | | | | | | | | |
| • | (Hybrid/Composite/Synthetic/Impr | | | | | | | | |
| oved/S11 | ngle Cross/Double Cross) | | | TT 1 1/10 1 | | | | | |
| S.No | Characteristics | States | Note | Hybrid/Research Variety | Female | Male | | | |
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | | | |
| | Plant: Growth habit | Erect | 1 | | | | | | |
| 1 | | Semi- spreading | 2 | | | | | | |
| | | Spreading | 3 | | | | | | |
| | Leaflet: Size (fully developed basal leaflet) | Small (<4.0 cm) | 3 | | | | | | |
| 2 | | Medium (<4.0 –6.0 cm) | 5 | | | | | | |
| | | Large (>6.0 cm) | 7 | | | | | | |
| | Leaflet: Colour | Light green | 1 | | | | | | |
| 3 | | Green | 2 | | | | | | |
| | | Dark green | 3 | | | | | | |
| 4 | Stem: Pubescence | Absent | 1 | | | | | | |
| | | Sparse | 3 | | | | | | |
| | | Medium | 5 | | | | | | |
| 5 | Flower: Presence on main axis | Absent | 1 | | | | | | |
| | | Present | 9 | | | | | | |
| 6 | Flower: Arrangement on side branches | Sequential | 1 | | | | | | |
| | | Alternate | 2 | | | | | | |
| | | Irregular | 3 | | | | | | |
| 7 | Inflorescence | Simple | 1 | | | | | | |
| | | Compound | 2 | | | | | | |
| 8 | Time of maturity (For curing) | Very early (< 90 days) | 1 | | | | | | |
| | | Early (90-100 days) | 3 | | | | | | |
| | | Medium (101-110 days) | 5 | | | | | | |
| | | Late (111-120 days) | 7 | | | | | | |
| | | Very late (>120 days) | 9 | | | | | | |
| 9 | Pod: Constriction | Absent | 1 | | | | | | |
| | | Shallow | 3 | | | | | | |
| | | Medium | 5 | | | | | | |
| | | Deep | 7 | | | | | | |
| 10 | Pod: Reticulation | Absent | 1 | | | | | | |
| | | Medium | 3 | | | | | | |
| | | Prominent | 5 | | | | | | |
| 11 | Pod: Number of kernels (on 100 pod basis) | >60% 2 seeded | 1 | | | | | | |
| | | >60% 3 seeded | 3 | | | | | | |
| | | >60% 4 seeded | 5 | | | | | | |
| 12 | Pod: Presence of beak | Absent | 1 | | | | | | |
| | | Present | 9 | | | | | | |
| | | Low (<66) | 3 | | | | | | |
| 13 | Pod: Shelling percentage | Medium (66-75) | 5 | | | | | | |
| | | High (>75) | 7 | 1 | 1 | | | | |

High (>75)

| S.No | Characteristics | States | Note | Hybrid/Research Variety | Female | Male |
|------|--|--------------------------|------|----------------------------|--------|------|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| 14 | Testa: Colour | Uniform | 1 | | | |
| | | Variegated | 9 | | | |
| | Kernel: Colour of testa (varieties with monochrome testa only) | White (1 A 1) | 1 | | | |
| | | Off white(1 A 2) | 2 | | | |
| | | Tan (12 E 4) | 3 |] | | |
| | | Rose (Grayish red 8 B 3) | 4 |] | | |
| 15 | | Purple (14 F 4) | 5 |] | | |
| | | Dark purple (14 F 7) | 6 | | | |
| | | Salmon (6 A 4) | 7 | | | |
| | | Red (10 B 7) | 8 | | | |
| | | Dark red (11 C 8) | 9 | | | |
| | Kernel: Shape | Spheroid | 1 | | | |
| 16 | | Cylindrical | 2 | | | |
| | | Fusiform | 3 | | | |
| 17 | Kernel: Weight of 100 kernels (about 9% moisture) | Low (<36 g) | 3 | | | |
| | | Medium (36-50 g) | 5 | | | |
| | | High (51-65 g) | 7 | | | |
| | | Very high (>65 g) | 9 | | | |
| 18 | Seed: Fresh seed dormancy | Absent | 1 | | | • |
| | | Present | 9 | | | |