Company	y Name							
Crop Name		SOYBEAN						
Variety N								
Types of								
	composite/ synthetic/ improved/							
Single cr	oss/ Double cross)							
S.No.	Characteristics	States	Note	Hybrid/Res earch	Female	Male		
1	2	3	4	5	6	7		
1	Hypocotyl: Anthocyanin	Absent	1					
1	pigmentation	Present	9	1				
	Plant: Growth type	Determinate	1					
2		Semi-determinate	2	1				
		Indeterminate	3	1				
	Plant: Days to 50% flowering	Early (≤ 35 days)	3					
3		Medium (36 to 45 days)	5	1				
		Late (> 45 days)	7					
		Lanceolate	1					
4	Leaf: Shape	Pointed ovate	2]				
		Rounded ovate	3					
5	Leaf : Colour	Green	1					
3		Dark green	2					
6	Plant: Growth habit	Erect	1					
O		Semi-erect	2					
7	Flower: Colour	White	1					
/		Purple	2					
	Plant: Height (cm)	Short (≤ 40)	3					
8		Medium (41-60)	5					
		Tall (>60)	7					
9	Pod: Pubescence	Absent	1					
		Present	9					
10	Pod: Pubescence colour	Grey	1					
10		Tawny (Brown)	2					
	Pod: Colour	Yellow	1	_				
11		Brown	2	_				
		Black	3					
12	Pod: Shattering	Shattering	1	1				
	r our sharrenng	Non shattering (up to 10-days)	9					
	Plant: Days to maturity	Early (≤ 95 days)	3	_				
13		Medium (96-105 days)	5	_				
		Late (>105 days)	7					
	Seed: Size (100 seeds weight)	Small (≤ 10.0 g)	3	1				
14		Medium (10.1-13.0 g)	5	_				
		Large (> 13.0 g)	7					
15	Seed: Shape	Spherical	1	4				
		Elliptical	2					
	Seed: Colour	Yellow	1	4				
16		Yellow green	2	4				
		Green	3	4				
		Black	4					
17	Seed: Lustre	Shiny	1	4				
		Dull	9					

S.No.	Characteristics	States	Note	Hybrid/Res earch	Female	Male
1	2	3	4	5	6	7
18	Seed: Colouration due to peroxidase activity in seed coat	Absent	1			
		Present	9	1		
19	Seed: Hilum colour	Yellow	1			
		Grey	2			
		Brown	3			
		Black	4			
		Variegated	5			
20	Seed: Cotyledon colour	Yellow	1			
		Green	2			
21	Seed: Oil content (%)	Low (≤ 15.0)	3			
		Average (15.1-18.0)	5			
		Medium (18.1-20.0)	7			
		High (> 20.0)	9			
22	Seed: Protein content (%)	Low (≤ 38.0)	3			
		Medium (38.1 – 40.0)	5			
		High (>40.0)	7			